
Curvature of the Universe


Jan J. Ostrowski

National Centre for Nuclear Research, Warsaw Annual seminar of Department of Fundamental Research, 15-12-2020



Spatial curvature in the FLRW universe


Cosmic triangle: $1=\Omega_m+\Omega_k+\Omega_\Lambda$



We think that these order–of–magnitude estimates provide a strong call for a proper relativistic treatment of the underlying gravitational physics in these systems; spatial curvature is an inherently relativistic phenomenon, unknown to the Newtonian theory. The claim on the validity of a quasi–Newtonian metric (...) to describe gravitational physics on all scales in the observable Universe (...) is thus seriously called into question.


Gravitating system / Smoothing scale	Mass M	Diameters D and d	D/d	ε	$\varepsilon (D/d)^2$
A1: Earth's orbit / Sun	$\begin{array}{c} \approx M_{\odot} \\ (1.99 \times 10^{30} \text{ kg}) \end{array}$	$\begin{array}{l} 300\times10^6~{\rm km}\\ 1.39\times10^6~{\rm km} \end{array}$	216	4.24×10^{-6}	0.20
A2: Galaxy / Open star cluster	$\approx 10^{11} M_{\odot}$ (1.99 × 10 ⁴¹ kg)	100000 ly 30 ly	3333	$6.23 imes 10^{-7}$	6.92
A3: Cluster of galaxies / Galaxy	$\approx 10^{14} M_{\odot}$ (1.99 × 10 ⁴⁴ kg)	5 Mpc 0.03 Mpc	167	3.82×10^{-6}	0.11
C1: Void / Wall	$\approx (1/6)\pi \rho_m D^3$ (2.98 × 10 ⁴⁵ kg)	$30h^{-1} \text{ Mpc} 3h^{-1} \text{ Mpc}$	10	6.78×10^{-6}	6.78×10^{-4}
C2: Homogeneity scale / Supercluster	$\approx (1/6)\pi\rho_m D^3$ $(2.98\times 10^{48} \text{ kg})$	$300h^{-1} \text{ Mpc}$ $30h^{-1} \text{ Mpc}$	10	6.78×10^{-4}	6.78×10^{-2}
C3: Hubble sphere /	$\approx (1/6)\pi\rho_m D^3$ $(2.38\times 10^{52} \text{ kg})$	$6000h^{-1}$ Mpc		0.27	

Spatial curvature: observational effect



Roukema, Ostrowski, Buchert; JCAP 2013

Partitioning approach



Buchert, Carfora; CQG 2002

Hamiltonian constraint, turnaround condition

Local Hamiltonian constraint:

$$3H^2 = 8\pi G\rho - 3k/a^2 + \Lambda \quad \Rightarrow \quad H^2 = 8\pi G\rho + \sigma^2 - \frac{1}{2}\mathcal{R} + \Lambda$$

Turnaround condition: H = 0

Hamiltonian constraint, turnaround condition

Local Hamiltonian constraint:

$$3H^2 = 8\pi G\rho - 3k/a^2 + \Lambda \quad \Rightarrow \quad H^2 = 8\pi G\rho + \sigma^2 - \frac{1}{2}\mathcal{R} + \Lambda$$

Turnaround condition: H = 0

Averaged Hamiltonian constraint:

$$H_{\mathcal{D}}^2 = 8\pi G \langle \rho \rangle_{\mathcal{D}} - \frac{1}{2} \mathcal{Q} - \frac{1}{2} \langle \mathcal{R} \rangle_{\mathcal{D}} + \Lambda$$

where Q contains kinematical effects from inhomogeneities

Turnaround condition: $H_{\mathcal{D}} = 0$

Analytical results

For the turnaround to occur:

 $\mathcal{R} > 0$

In the case of averaged equations, we have statistically:

 $\langle \mathcal{R} \rangle_{\mathcal{D}} > 0$

For Einstein de-Sitter background we have:


$$\Omega_{\mathcal{R}}^{\mathcal{D}} = -5 \; ; \; \Omega_{\mathcal{Q}}^{\mathcal{D}} = 1 \; ; \; \Omega_{m}^{\mathcal{D}} = 4 \; ; \; \frac{\langle \rho \rangle_{\mathcal{D}}}{\rho_{EdS}} = 4$$

Numerical methods


General scheme:

- **MPGRAFIC** generate initial conditions
- DTFE calculate averaged initial conditions
- **INHOMOG** calculate evolution of the domains
- **RAMSES-SCALAV** single pipeline + additional options

Curvature density



Averaged positive curvature



Conclusions

- big positive spatial curvature is a generic feature of collapsing structures at the turnaround; both locally and on average
- $\Omega_R^D \approx -5$ remains an approximate lower bound for the averaged curvature functional for the wide range of initial conditions
- fluid parameters at the turnaround may provide an additional cosmological test
- details can be found in:
 - \rightarrow 'A few numbers from the turnaround epoch of collapse', Ostrowski J.J., Acta Phys. Pol. B Proc. Suppl. Vol. 13, p 177 (2020)
 - → 'Does spatial flatness forbid the turnaround epoch of collapsing structures?',
 Roukema B.F., Ostrowski J.J., Journal of Cosmology and Astroparticle Physics, Vol. 2019, 12 (2019)