

Hunting for Red nuggets untouched survivors of the early Universe

Krzysztof Lisiecki

Katarzyna Małek

Agnieszka Pollo

Junais



VIMOS PUBLIC EXTRAGALACTIC REDSHIFT SURVEY




NATIONAL CENTRE
FOR NUCLEAR RESEARCH
ŚWIERK

What red nuggets really are?



VIPERS red nuggets
Krzysztof Lisiecki

Red nuggets are a rare population of passive compact massive galaxies thought to be the first massive galaxies that formed in the Universe.




What red nuggets really are?




VIPERS red nuggets
Krzysztof Lisiecki

But the mergers events are stochastic,
so we can expect some of them in the
local or, at least, in the closer
Universe.



Why those are important?

Discovery of massive, compact and passive objects at high-redshift challenged the cosmological models. As a response to this problem, two-phase scenario was proposed.




Red nuggets: how to find them?



VIPERS red nuggets
Krzysztof Lisiecki

There are three things that determine
red nugget:


1. stellar mass higher than 10^{10} – 10^{11} solar masses
2. size smaller than a few kpc
3. low star formation rate – passiveness



Red nugget
Milky Way





- ~90k spectroscopically measured galaxies;
- redshift range: 0.4 - 1.2;
- wavelength range: 450 - 950 nm;
- total area: 23.5 deg²;




To derive physical properties, in particular stellar masses and SFRs, we used the Code Investigating The Galaxy Emission (CIGALE).

The morphological parameters were derived by Krywult et. al 2017.



Almighty VIPERS red nuggets selection: preselection

- 95% confidence in redshift estimation to ensure effective radii in kpc
- $0.5 < z < 1$ to be complete in colour




Fritz et al. 2014

| Cut | Sample size |
|------------------------------------|-------------|
| VIPERS database | 91 507 |
| $z_{flag} \in \{3, 4, 23, 24\}$ | 54 252 |
| Redshift range $0.5 \leq z \leq 1$ | 44 145 |
| R_e uncertainties | 36 157 |

Almighty VIPERS red nuggets selection: compactness

As we wanted to be sure that we select only truly compact and massive sources, we decided to use one of the most restrictive criterion (Trujillo et. al 2009):

| Reference | Number of sources |
|--|-------------------|
| Damjanov et al. (2015) | 4 347 |
| Cassata et al. (2011) – compact | 3 139 |
| Barro et al. (2013) | 3 083 |
| van der Wel et al. (2014) – compact | 1 801 |
| Charbonnier et al. (2017) | 1 061 |
| Spiniello et al. (2021) | 693 |
| Buitrago et al. (2018) | 277 |
| Cassata et al. (2011) – ultracompact | 250 |
| van der Wel et al. (2014) – ultracompact | 241 |
| Trujillo et al. (2009) | 86 |




$M_{\text{star}} > 8 \times 10^{10} M_{\odot}$ and $R_e < 1.5 \text{ kpc}$

Almighty VIPERS red nuggets selection: passiveness



We performed multistage selection based on colours, emission lines, and visual check.



NUVrK diagram is widely used by VIPERS team to separate red and blue galaxy populations.



Almighty VIPERS red nuggets selection: passiveness II

Passive/active spectra

Main sequence



Almighty VIPERS red nuggets catalogue




VIPERS red nuggets
Krzysztof Lisiecki

We established the first spectroscopic catalogue of red nuggets at z 0.5-1. In total 77 sources, which is the largest spec- z sample above $z \sim 0.5$.

Divided them into three redshift bins:

| Redshift range | N |
|-------------------------|----|
| $0.50 \leq z \leq 0.75$ | 22 |
| $0.75 < z \leq 0.90$ | 33 |
| $0.90 < z \leq 1.00$ | 22 |
| $0.50 \leq z \leq 1.00$ | 77 |




Almighty VIPERS red nuggets catalogue: number densities

Calculated number densities per cubic comoving Mpc are in agreement with low- z measurements but not so much with high- z .



| Reference | Number of sources |
|--|-------------------|
| Barro et al. (2013) | 3 083 |
| van der Wel et al. (2014) – ultracompact | 241 |
| Tortora et al. (2016) | 86 |

| Redshift range | Number density [Mpc^{-3}] |
|-------------------------|--------------------------------------|
| $0.50 \leq z \leq 0.75$ | 5.46×10^{-6} |
| $0.75 < z \leq 0.90$ | 1.22×10^{-5} |
| $0.90 < z \leq 1.00$ | 9.82×10^{-6} |
| $0.50 \leq z \leq 1.00$ | 8.86×10^{-6} |





To characterize the environments of red nuggets,
we generated three control samples.

All red
VIPERS
galaxies



Similar mass
larger size

Similar size
smaller mass




Red nuggets



Red nuggets do not have environmental preferences.


We found 11 red nuggets in low density and 10 in high density environment.

The most similar distribution can be found in sample with galaxies with similar sizes and lower masses.



Summary

- We found 77 spectroscopically selected red nuggets at intermediate redshift. It is the first catalogue of this kind.
- All of them are spectroscopically identified – unique for red nuggets
- Number densities are in good agreement – it is not trivial to compare due to the selection function
- We found no relation with the environment.



The established catalogue is just a beginning and a perfect starting sample for future studies:

1. studying stellar populations;
2. looking for relics;
3. studying individual galaxies;
4. studying statistics;
5. focusing on compact, but not so passive sources;
6. and many more...

Thank you for your attention!



This projects is supported by Ministry of Education and Science
via grant PN/01/0034/2022!

Almighty VIPERS: photometry

VIPERS red nuggets
Krzysztof Lisiecki

| Telescope/ Instrument | Filter | λ_{mean} (μm) |
|--------------------------|------------------|---------------------------------------|
| GALEX | FUV | 0.155 |
| | NUV | 0.234 |
| CFHT/MegaCam | <i>u</i> | 0.369 |
| | <i>g</i> | 0.482 |
| | <i>r</i> | 0.643 |
| | <i>i</i> | 0.772 |
| | <i>z</i> | 0.900 |
| | <i>iy</i> | 0.769 |
| | CFHT/Wircam | K_s |
| VISTA | K_{video} | 2.158 |
| WISE | W1 | 3.353 |
| | W2 | 4.603 |
| | W3 | 11.561 |
| | W4 | 22.088 |
| | Spitzer/IRAC | I1 |
| Spitzer/MIPS | I2 | 4.505 |
| | I3 | 5.739 |
| | I4 | 7.927 |
| | 24 μm | 23.843 |
| 70 μm | 72.555 | |
| 160 μm | 157.000 | |





Spectroscopic redshifts with >95% confidence for 54 252 galaxies

Spectra restframe wavelength range:

$z = 0.5 \rightarrow 3000 - 6300 \text{ \AA}$

$z = 1.0 \rightarrow 2250 - 4750 \text{ \AA}$



Half-light radii, θ_e , was derived using GALFIT with Sersic profile:

$$I(r) = I_e \exp \left(-b_n \left[\left(\frac{r}{\theta_e} \right)^{1/n} - 1 \right] \right)$$

In analysis we used circuralised half-light radii:

$$R_e = \theta_e \sqrt{b/a}$$

